




What you'll Learn About

- Right Triangle Trigonometry/ Two Famous Triangles
- Evaluating Trig Functions with a calculator/Applications of right triangle trig

The six trigonometric functions

Find the values of all six trigonometric functions.



Assume that  $\theta$  is an acute angle in a right triangle satisfying the given conditions. Evaluate the remaining trigonometric functions.

A)  $\sin \theta = \frac{4}{9}$

B)  $\cos \theta = \frac{2}{9}$


C)  $\tan \theta = \frac{4}{9}$

D)  $\cot \theta = \frac{2}{9}$


E)  $\csc \theta = \frac{10}{7}$

F)  $\sec \theta = \frac{4}{3}$

## 45-45-90 Triangle



## 30-60-90 Triangle



Evaluate using a calculator. Make sure your calculator is in the correct mode. Give answers to 3 decimal places and then draw the triangle that represents the situation.

A)  $\sin 53^\circ$

B)  $\cos \frac{2\pi}{5}$

C)  $\tan 154^\circ$


D)  $\cot \frac{\pi}{9}$

E)  $\csc 220^\circ$


F)  $\sec \frac{8\pi}{5}$

Solve the triangle for the variable shown.

9)



10)



Solve the triangle ABC for all of its unknown parts. Assume C is the right angle.

$$\alpha = 40^\circ \quad a = 10$$

Solve the triangle ABC for all of its unknown parts. Assume C is the right angle.

$$\beta = 62^\circ \quad a = 7$$

Example 6: From a point 340 feet away from the base of the Peachtree Center Plaza in Atlanta, Georgia, the angle of elevation to the top of the building is  $65^\circ$ . Find the height of the building.



--	--

What you'll Learn About

- Trig functions of any angle/Trig functions of real numbers
- Periodic Functions/The Unit Circle

Point P is on the terminal side of angle  $\theta$ . Evaluate the six trigonometric functions for  $\theta$ .

A) (5, 4)

B) (-3, 4)

C) (-2, -5)

D) (-4, -1)

E) (0, -3)

F) (3, 0)

Determine the sign (+ or -) of the given value without the use of a calculator.

A)  $\sin 53^\circ$

B)  $\cos \frac{2\pi}{5}$


C)  $\tan 154^\circ$

D)  $\cot \frac{\pi}{9}$

E)  $\csc 220^\circ$

F)  $\sec \frac{8\pi}{5}$

# Unit Circle, Fill in the blank



Evaluate without using a calculator by using ratios in a reference triangle.

A)  $\sin 120^\circ$

B)  $\cos \frac{2\pi}{3}$

C)  $\tan \frac{13\pi}{4}$

D)  $\cot \frac{-13\pi}{6}$

E)  $\csc \frac{7\pi}{4}$

F)  $\sec \frac{23\pi}{6}$

Find sine, cosine, and tangent for the given angle.

A)  $90^\circ$

B)  $-\frac{\pi}{2}$

C)  $6\pi$

D)  $\frac{-7\pi}{2}$

Evaluate without using a calculator

A) Find  $\sin \theta$  and  $\tan \theta$  if  $\cos \theta = \frac{3}{4}$  and  $\cot \theta < 0$


B) Find  $\sec \theta$  and  $\csc \theta$  if  $\cot \theta = \frac{-6}{5}$  and  $\sin \theta > 0$

--	--


What you'll Learn About

- The basic waves revisited/Sinusoids and Transformations
- Modeling

The graph of  $y = \sin x$



The graph of  $y = \cos x$





Find the amplitude of the function and use the language of transformations to describe how the graph of the function is related to the graph of  $y = \sin x$

A)  $y = 3\sin x$

B)  $y = \frac{3}{4}\sin x$

C)  $y = -5\sin x$

Find the period of the function and use the language of transformations to describe how the graph of the function is related to the graph of  $y = \cos x$

A)  $y = \cos(2x)$

B)  $y = \cos \frac{x}{2}$

C)  $y = \cos\left(\frac{-3x}{4}\right)$

Graph 1 period of the function without using your calculator.

A)  $y = 3 \sin \frac{x}{2}$

$$y = 5 \cos 2x$$

Identify the maximum and minimum values and the zeros of the function in the interval  $[-2\pi, 2\pi]$ . Use your understanding of transformations, not your calculator.

A)  $y = 4 \sin x$

B)  $y = -2 \cos \frac{x}{3}$

Determine the phase shift for the function and the sketch the graph.

A)  $y = \cos\left(x - \frac{\pi}{6}\right)$

B)  $y = \sin\left(x + \frac{\pi}{3}\right)$

Determine the vertical shift for the function and the sketch the graph.

A)  $y = \cos x - 2$

B)  $y = \sin x + 3$

Determine the vertical shift and phase shift of the function and then sketch the graph

A)  $y = \cos\left(x + \frac{\pi}{6}\right) - 1$

B)  $y = \sin\left(x - \frac{\pi}{3}\right) + 2$

State the Amplitude and period of the sinusoid, and relative to the basic function, the phase shift and vertical translation.

A)  $y = 3\sin\left(x - \frac{\pi}{4}\right) + 2$

B)  $y = -2\cos\left(3x - \frac{\pi}{4}\right) - 4$

C)  $y = 5\sin 4\pi x + 6$

$$\text{Amp} = A = \frac{\text{Max} - \text{Min}}{2}$$

$$\text{Vertical} = (C) = \frac{\text{Max} + \text{Min}}{2}$$

$$\text{period} = p$$

Horizontal Stretch/Shrink

$$B = \frac{2\pi}{p}$$

How to choose an appropriate model based on the behavior at some given time, T.

$y = A \cos B(t - T) + C$   
if at time T the function attains a maximum value

$y = -A \cos B(t - T) + C$   
if at time T the function attains a minimum value

$y = A \sin B(t - T) + C$   
if at time T the function halfway between a minimum and a maximum value

$y = -A \sin B(t - T) + C$   
if at time T the function halfway between a maximum and a minimum value

Construct a sinusoid with the given amplitude and period that goes through the given point.

A) Amp: 4, period  $4\pi$ , point (0, 0)

B) Amp: 2.5, period  $\frac{\pi}{5}$ , point (2, 0)

$$\text{Amp} = A = \frac{\text{Max} - \text{Min}}{2}$$

$$\text{Vertical} = (C) = \frac{\text{Max} + \text{Min}}{2}$$

$$\text{period} = p$$

Horizontal Stretch/Shrink

$$B = \frac{2\pi}{p}$$

How to choose an appropriate model based on the behavior at some given time, T.

$y = A \cos B(t - T) + C$   
if at time T the function attains a maximum value

$y = -A \cos B(t - T) + C$   
if at time T the function attains a minimum value

$y = A \sin B(t - T) + C$   
if at time T the function halfway between a minimum and a maximum value

$y = -A \sin B(t - T) + C$   
if at time T the function halfway between a maximum and a minimum value

### Example 7: Calculating the Ebb and Flow of Tides

One particular July 4<sup>th</sup> in Galveston, TX, high tide occurred at 9:36 am. At that time the water at the end of the 61<sup>st</sup> Street Pier was 2.7 meters deep. Low tide occurred at 3:48 p.m, at which time the water was only 2.1 meters deep. Assume that the depth of the water is a sinusoidal function of time with a period of half a lunar day (about 12 hrs 24 min)

a) Model the depth, D, as a sinusoidal function of time, t, algebraically then graph the function.

b) At what time on the 4<sup>th</sup> of July did the first low tide occur.

c) What was the approximate depth of the water at 6:00 am and at 3:00 pm?

d) What was the first time on July 4<sup>th</sup> when the water was 2.4 meters deep?

80) Temperature Data: The normal monthly Fahrenheit temperatures in Helena, MT, are shown in the table below (month 1 = January)

Model the temperature  $T$  as a sinusoidal function of time using 20 as the minimum value and 68 as the maximum value. Support your answer graphically by graphing your function with a scatter plot.

M	1	2	3	4	5	6	7	8	9	10	11	12
T	20	26	35	44	53	61	68	67	56	45	31	21


--	--




What you'll Learn About

- The graphs of the other 4 trig functions


The graph of  $y = \csc x$




The graph of  $y = \sec x$



The graph of  $y = \tan x$



The graph of  $y = \cot x$



Describe the graph of the function in terms of a basic trigonometric function. Locate the vertical asymptotes and graph 2 periods of the function.

A)  $y = 2\tan(3x)$

B)  $y = -\cot(2x)$

C)  $y = \sec(4x)$

D)  $y = -\csc\left(\frac{x}{3}\right)$

Describe the transformations required to obtain the graph of the given function from a basic trigonometric graph.

A)  $y = 5 \tan x$

B)  $y = -3 \cot\left(\frac{x}{2}\right)$

C)  $y = 2 \sec \frac{4x}{3}$


D)  $y = -4 \csc 2\pi x - 3$

--	--


What you'll Learn About

- Inverse Trigonometric Functions and their Graphs


The graph of  $y = \sin x$




The graph of  $y = \sin^{-1} x = \arcsin x$




# The Unit Circle and Inverse Functions




The graph of  $y = \cos x$



The graph of  $y = \cos^{-1} x = \arccos x$




The Unit Circle and Inverse Functions







The graph of  $y = \tan x$



The graph of  $y = \tan^{-1} x = \arctan x$



Find the exact value

A)  $\cos^{-1} \frac{\sqrt{3}}{2}$

B)  $\cos^{-1} \frac{1}{2}$

C)  $\cos^{-1} \left( \frac{-1}{2} \right)$

D)  $\sin^{-1} \frac{-\sqrt{3}}{2}$

E)  $\sin^{-1} \frac{1}{2}$

F)  $\sin^{-1} \left( \frac{1}{\sqrt{2}} \right)$

G)  $\tan^{-1}(1)$

H)  $\tan^{-1}(\sqrt{3})$

I)  $\tan^{-1} \left( \frac{-1}{\sqrt{3}} \right)$

J)  $\cos^{-1}(0)$

K)  $\sin^{-1}(-1)$

L)  $\tan^{-1}(0)$

Use a calculator to find the approximate value in degrees. Draw the triangle that represents the situation.

A)  $\arccos(.456)$

B)  $\arcsin(-.456)$

C)  $\arctan(-5.768)$

Use a calculator to find the approximate value in radians. Draw the triangle that represents the situation.

A)  $\arcsin(.456)$

B)  $\arccos(-.456)$

C)  $\arctan(-5.768)$

Find the exact value without a calculator.

A)  $\sin(\cos^{-1}(1/2))$

B)  $\cos(\tan^{-1}(0))$

C)  $\tan\left(\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)\right)$

D)  $\sin(\tan^{-1}(-\sqrt{3}))$

E)  $\cos^{-1}\left(\sin\left(\frac{\pi}{4}\right)\right)$

F)  $\sin^{-1}\left(\cos\left(\frac{\pi}{6}\right)\right)$

--	--

*PRE-CALCULUS: by Finney, Demana, Watts and Kennedy*  
*Solving Trigonometric Equations*

What you'll Learn About

Solve each trigonometric equation for  $\theta$  on the interval  $[0, 2\pi]$ . Then give a formula for all possible angles that could be a solution of the equation.

A)  $\sin \theta = \frac{\sqrt{2}}{2}$

B)  $\cos \theta = \frac{-1}{2}$

C)  $\sin \theta = 1$

D)  $\cos \theta = 0$

E)  $\tan \theta = \sqrt{3}$

F)  $\tan \theta = -1$

Solve each trigonometric equation for  $\theta$  on the interval  $[0, 2\pi]$ .

A)  $\cos 2\theta = \frac{1}{2}$

B)  $\sin 3\theta = \frac{1}{2}$

C)  $\cos \frac{\theta}{3} = \frac{\sqrt{3}}{2}$

D)  $\tan\left(\frac{\theta}{2} + \frac{\pi}{3}\right) = 1$

E)  $\sin \theta = .4$

F)  $\cos \theta = -.2$

$$\text{A) } \sqrt{2} \cos \theta - 1 = 0$$

$$\text{B) } \sqrt{3} \csc \theta - 2 = 0$$

$$\text{C) } 4 \sin^2 \theta - 1 = 0$$

$$\text{D) } (3 \cot^2 \theta - 1)(\cot^2 \theta - 3) = 0$$

$$\text{E) } 3 \tan^2 \theta - 1 = 0$$

$$\text{F) } \cos^2 \theta = 3 \sin^2 \theta$$



$$\text{G) } 2\cos^2 \theta + \cos \theta = 0$$

$$\text{H) } 2\sin \theta \cos \theta = \cos \theta$$

$$\text{I) } \csc^2 \theta - \csc \theta = 2$$

$$\text{J) } \sin^3 \theta = \sin \theta$$

--	--

**Reciprocal Identities**

$$\begin{aligned} \csc x &= \frac{1}{\sin x} & \sec x &= \frac{1}{\cos x} & \cot x &= \frac{1}{\tan x} \\ \sin x &= \frac{1}{\csc x} & \cos x &= \frac{1}{\sec x} & \tan x &= \frac{1}{\cot x} \end{aligned}$$

**Quotient Identities**

$$\tan x = \frac{\sin x}{\cos x} \quad \cot x = \frac{\cos x}{\sin x}$$

**Even/Odd Identities**

$$\begin{aligned} \sin(-x) &= -\sin x & \csc(-x) &= -\csc x \\ \cos(-x) &= \cos x & \sec(-x) &= \sec x \\ \tan(-x) &= -\tan x & \cot(-x) &= -\cot x \end{aligned}$$

**Pythagorean Identities**

$$\begin{aligned} \sin^2 x + \cos^2 x &= 1 & \tan^2 x + 1 &= \sec^2 x & 1 + \cot^2 x &= \csc^2 x \\ \sin^2 x &= 1 - \cos^2 x & \tan^2 x &= \sec^2 x - 1 & \csc^2 x - \cot^2 x &= 1 \\ \cos^2 x &= 1 - \sin^2 x & \sec^2 x - \tan^2 x &= 1 & \cot^2 x &= \csc^2 x - 1 \end{aligned}$$

**Co-function**

$$\begin{aligned} \sin\left(\frac{\pi}{2} - x\right) &= \cos x & \cos\left(\frac{\pi}{2} - x\right) &= \sin x & \tan\left(\frac{\pi}{2} - x\right) &= \cot x \\ \csc\left(\frac{\pi}{2} - x\right) &= \sec x & \sec\left(\frac{\pi}{2} - x\right) &= \csc x & \cot\left(\frac{\pi}{2} - x\right) &= \tan x \end{aligned}$$

Equation of Unit  
Circle

$$x^2 + y^2 = 1$$

Use trig ratios to prove that  $\cos^2 \theta + \sin^2 \theta = 1$   
is the equation of the unit circle

Use basic identities to simplify the expression to a different trig function  
or a product of two trig functions

10.  $\cot x \tan x$

A.  $\frac{1 - \sin^2 \theta}{\cos \theta}$

B.  $\sin x - \sin^3 x$

C.  $\frac{\sin^2 x + \cot^2 x + \cos^2 x}{\csc x}$

Simplify the expression to either 1 or -1

17.  $\sin x \csc(-x)$

19.  $\cot(-x)\cot\left(\frac{\pi}{2}-x\right)$

21.  $\sin^2(-x)+\cos^2(-x)$

Simplify the expression to either a constant or a basic trig function.

A)  $\frac{\cot\left(\frac{\pi}{2}-x\right)\sec x}{\sec^2 x}$

B)  $\frac{1+\cot x}{1+\tan x}$

C)  $\tan^2 x + \cot^2 x - (\sec^2 x + \csc^2 x)$

Use the basic identities to change the expression to one involving only sines and cosines. Then simplify to a basic trig function.

$$28) \sin \theta - \tan \theta \cos \theta + \cos\left(\frac{\pi}{2} - \theta\right)$$

$$30) \frac{(\sec y - \tan y)(\sec y + \tan y)}{\sec y}$$

$$31. \frac{\tan x}{\csc^2 x} + \frac{\tan x}{\sec^2 x}$$

Combine the fractions and simplify to a multiple of a power of a basic trig function

A) 
$$\frac{1}{1 - \cos x} + \frac{1}{1 + \cos x}$$

35. 
$$\frac{\sin x}{\cot^2 x} - \frac{\sin x}{\cos^2 x}$$

Write each expression in factored form as an algebraic expression of a single trig function

A)  $\sin^2 x + 2\sin x + 1$

B)  $1 - 2\cos x + \cos^2 x$

C)  $\sin x - 2\cos^2 x + 1$

45.  $4\tan^2 x - \frac{4}{\cot x} + \sin x \csc x$

Write each expression as an algebraic expression of a single trigonometric function

A)  $\frac{1 - \cos^2 x}{1 + \cos x}$

B)  $\frac{\cot^2 \alpha - 1}{1 + \cot x}$

C)  $\frac{\cos^2 x}{1 + \sin x}$

D)  $\frac{\cot^2 x}{1 + \csc x}$



What you'll Learn About

$$12. \sin x(\cot x + \cos x \tan x) = \cos x + \sin^2 x$$

$$14. (\cos x - \sin x)^2 = 1 - 2 \sin x \cos x$$

$$18. \frac{\sec^2 \theta - 1}{\sin \theta} = \frac{\sin \theta}{1 - \sin^2 \theta}$$

$$20. \frac{1}{1 - \cos x} + \frac{1}{1 + \cos x} = 2 \csc^2 x$$

$$22. \sin^2 \alpha - \cos^2 \alpha = 1 - 2\cos^2 \alpha$$

$$26. \frac{\sec x + 1}{\tan x} = \frac{\sin x}{1 - \cos x}$$

$$30. \quad \tan^2 \theta - \sin^2 \theta = \tan^2 \theta \sin^2 \theta$$

$$35. \quad \frac{\tan x}{\sec x - 1} = \frac{\sec x + 1}{\tan x}$$

$$37. \quad \frac{\sin x - \cos x}{\sin x + \cos x} = \frac{2 \sin^2 x - 1}{1 + 2 \sin x \cos x}$$

--	--

